

**POLYNOMIAL TIME LOGIC:
INABILITY TO EXPRESS
SH634**

SAHARON SHELAH

Institute of Mathematics
The Hebrew University
Jerusalem, Israel

Rutgers University
Mathematics Department
New Brunswick, NJ USA

Partially supported by the United States-Israel Binational Science Foundation
I would like to thank Alice Leonhardt for the beautiful typing.
Done - Fall 1996
Latest Revision - 98/July/23

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

ANOTATED CONTENT

- §1 The polynomial time logic presented
[We present this logic from a paper of Blass, Gurevich, Shelah [BGSh 533]; to compute what you can compute from a model in polynomial time without arbitrary choices (like ordering the model).]
- §2 The general elimination of quantifiers and proof it's non-expressive
[We define a criterion for showing the logic cannot say too complicated things on some model using a family of partial automorphism (rather than real automorphisms) and prove that it works. This is a relative of the Ehrenfeucht-Fraisse games, and more recent pebble games.]
- §3 The canonical example
[We deal with random enough graphs and conclude that they satisfy the 0-1 law so proving the logic cannot express two strong properties.]
- §4 Closing comments
[We present a variant of the criterion (the existence of a simple k -system). We then define a logic which naturally expresses it. We comment on defining $N_t[M]$ for ordinals.]

§1 THE POLYNOMIAL TIME LOGIC PRESENTED

We present below the choiceless polynomial time logic, introduced under the name $\tilde{CPTime}$ in Blass Gurevich Shelah [BGSh 533]. Knowledge of [BGSh 533] which is phrased with ASM (abstract state machine) is not required except when we explain how the definitions fit in 1.3(4). See on more relevant works there. The aim of this logic is to capture statements on a (finite) model M in polynomial time and space without arbitrary choices but with no additional bound on the depth, so its being this logic is a thesis. So we are not allowed to use a linear order on M , but if P^M has $\log_2(\|M\|)$ elements we are allowed to list all subsets of P^M , and if e.g. $(|P^M|)! \leq \|M\|$ we can list the permutations of P^M . Formally for a given M , we consider the elements of M as urelements, and build inductively $N_t = N_t[M]$, with $N_0 = M$, $N_{n+1} \subseteq N_t[M] \cup \mathcal{P}(N_t[M])$ but the definition is uniform and $N_t[M]$ should not be too large (i.e. has a polynomial bound) and the process stops.

Informally, we start with a model M with each element an atom=ure-element, we successively define $N_t[M]$, t running on the stages of the “computations”; to $N_{t+1}[M]$ we add few families of subsets of N_t , each of those defined by a $\psi(-, \bar{a})$ -formula for some $\bar{a}$ from $N_t[M]$, and we update few relations or functions, by defining them from those of the previous stage. Those are coded by $c_{t,\ell}$. We may then check if a target condition holds, then finishing. Note that each stage increases the size of $N_t[M]$ at most by a (fix) power, but in $\|M\|$ steps we can arrive to a model of size $2^{\|M\|}$. So we shall have a timing function $\mathbf{t}$ in $\|M\|$, normal polynomial, so when we have wasted too many resources (e.g. $\|N_t[M]\| + t$) our time is up whether we got an answer or not.

More formally

1.1 Definition. 1) For a model M , with vocabulary $\tau = \tau_0, \tau$ finite and $\in$ not in τ , let $\tau^+ = \tau_1 = \tau \cup \{\in\}$ considering the elements of M as atoms = urelements, we define $V_t[M]$ by induction of $t : V_0[M] = (M, \in \upharpoonright M)$ with $\in \upharpoonright M$ being empty (as we consider the members of M as atoms = “urelements”). Next $V_{t+1}[M]$ is the model with universe $V_t[M] \cup \{a : a \subseteq V_t[M]\}$ (so we assume $a \subseteq V_t[M] \Rightarrow a \notin M$ by “urelements”) with the predicates and individual constants and function symbols of τ interpreted as in M (so function symbols in τ are interpreted as partial functions) and $\in^{V_{t+1}[M]}$ is $\in \upharpoonright V_{t+1}[M]$.

2) We say $\Upsilon = (\bar{\psi}, \bar{\varphi}, \bar{c})$ is an inductive scheme for the language $\mathcal{L}_{f.o.}(\tau^+)$ (where $\mathcal{L}_{f.o.}$ is first order logic) if: letting $m_0 = \lg(\bar{\psi}), m_1 = \lg(\bar{\varphi})$ we have

- (a) $\bar{\mathcal{P}} = \langle \mathcal{P}_\ell : \ell < m_1 \rangle$ is a sequence of unary¹ predicates not in τ^+
- (b) $\bar{\psi} = \langle \psi_\ell : \ell < m_0 \rangle, \psi_\ell = \psi_\ell(x; \bar{y}_\ell)$ is first order in the vocabulary $\tau_2 = \tau_2[\Upsilon] = \tau_1 \cup \{\mathcal{P}_\ell : \ell < \lg(\bar{c})\}$
- (c) $\bar{\varphi} = \langle \varphi_\ell : \ell < m_1 \rangle, \varphi_\ell = \varphi_\ell(x)$ is first order in the vocabulary τ_2 .

3) For $M, \tau = \tau_0, \tau_1, \Upsilon = (\bar{\psi}, \bar{\varphi}, \bar{c})$ and τ_2 as above, we define by induction on t a submodel $N_t = N_t[M] = N_{\Upsilon,t}[M]$ of $V_t[M]$ and $\bar{c}_t = \langle c_{t,\ell} : \ell < m_1 \rangle$ with

¹but they will usually be interpreted as singletons, i.e. acts as individual constants but for the crucial “times” $t = 0, 1$ not necessarily

$c_{t,\ell} \in N_{t+1}, N_t^+[M]$ and $P_{t,\ell}, \mathcal{P}_{t,k}$ (for $\ell < m_0, k < m_1$) as follows: more exactly $N_t[M, \Upsilon], c_{t,0}[M, \Upsilon], \bar{c}_t[M, \Upsilon], P_{t,\ell}[M, \Upsilon], \mathcal{P}_{t,k}[M, \Upsilon]$.

We define $N_t[M]$ and $\mathcal{P}_{s,\ell}, c_{s,\ell}$ for $s < t$ by induction on t .

Case 1 $t = 0$: $N_t[M] = V_0[M]$.

Case 2 $t + 1$: $N_{t+1}[M]$ is the submodel of $V_{t+1}[M]$ with set of elements $N_t[M] \cup \bigcup_{\ell < m_0} \mathcal{P}_{t,\ell}[M]$

where

We define $\mathcal{P}_{t,k}[M], N_t^+[M]$ and $c_{t,\ell}$ by:

$$\mathcal{P}_{t,k}[M] = \left\{ \{a \in N_t[M] : N_t^+[M] \models \psi_k(a, \bar{b})\} : \bar{b} \in {}^{(\ell g(\bar{y}_i))}(N_t[M]) \right\}$$

$$P_{t,\ell}[M] = \{c_{s,\ell} : s < t \text{ and } s \text{ is the immediate successor of } s\}$$

$$N_t^+[M] = (N_t[M], P_{t,0}[M], \dots, P_{t,m-1}[M])$$

$$c_{t,\ell} = \{a \in N_t[M] : N_t^+[M] \models \varphi_\ell[a]\}$$

(so for $t > 1$, $\psi_k(x, \bar{y})$ is actually $\psi'_k(x, \bar{y}, \bar{c}_{t-1}) \in \mathcal{L}_{f.o.}(\tau_1)$).

Case 3: $t = \infty$ (i.e. w)

$$N_t = \bigcup_{s \leq t} N_s.$$

4) We say Υ is standard if some ψ_i guarantees that $\{s : s < t\} \subseteq N_t[M]$ (remember that we identify the natural number t with the set $\{0, 1, \dots, t-1\}$).

5) Let $\text{q.d.}(\varphi)$ be the quantifier depth of the formula φ . Let $m_{\text{qd}}(\Upsilon) = \text{Max}\{\text{q.d.}(\varphi) : \varphi \in \{\varphi_k : k < m_1\} \cup \{\psi_\ell : \ell < m_0\}\}$ and $m_{\text{fv}}(\Upsilon) = \text{Max}\{\ell g(\bar{y}_\ell) : \ell < m_0\}, \bar{m}(\Upsilon) = (m_{\text{qd}}(\Upsilon), m_{\text{fv}}(\Upsilon))$.

6) We may replace above first order logic by a logic $\mathcal{L}$. We let $\mathcal{L}_{f.o.}$ for first order, $\mathcal{L}_{\text{card}}$ like first order but we demand that Υ is standard and we allow the formulas $|\{x : \theta(x, \bar{y})\}| = s$. We let $\mathcal{L}_{\text{card}, \mathbf{T}}$ (on $\mathbf{T}$ see below) be like $\mathcal{L}_{f.o.}$ but for each $\mathbf{t} \in T$ we allow the quantifier $(Q_{\mathbf{t}}x)\varphi(x, \bar{y})$ with $N \models (Q_{\mathbf{t}}x)\varphi(x, \bar{a})$ iff $\mathbf{t}(\text{ure}(N)) < |\{b : N \models \varphi(b, \bar{a})\}|$ where $\text{ure}(N)$ is the set of urelements of N .

In the definition below the reader can concentrate on $\iota = 2$.

1.2 Definition. Let $\mathbf{T}$ be a set of functions $\mathbf{t} : \mathbb{N} \rightarrow \mathbb{N} \cup \{\infty\}$ and $\mathcal{L}^*$ be a logic ($\mathcal{L}_{f.o.}$ or $\mathcal{L}_{\text{card}}$ usually) and let τ be a vocabulary. If $\mathbf{t}$ is constantly ∞ we may write ∞ .

We define for $\iota = 1, 2, 3, 4$ the logic $\mathcal{L}_\iota^{\mathbf{T}}[\mathcal{L}^*](\tau)$, for all of those logics the set of sentences for a vocabulary τ is a subset of $\Theta = \Theta_\tau = \{\theta_{\Upsilon, \chi, \mathbf{t}} : \Upsilon \text{ an inductive scheme for } \mathcal{L}^*(\tau), \chi \in \mathcal{L}^*(\tau) \text{ and } \mathbf{t} \in \mathbf{T}\}$, (equal if not said otherwise), and for most of them we define the stopping time $t_\iota(M, \Upsilon, \mathbf{t})$ or $t_\iota(M, \Upsilon)$ (if $\mathbf{t}$ does not matter). The satisfaction relation for $\mathcal{L}_\iota^{\mathbf{T}}[\mathcal{L}^*](\tau)$ is denoted by $\models_\iota$. Also we write $\theta_{\Upsilon, \chi}$ if $\mathbf{t}$ does not matter. We may let $\text{Dom}(\mathbf{t})$ be the set of relevant structures.

Case 1: For $\iota = 1$.

$$t_\iota[M, \Upsilon] = \text{Min}\{t : t \geq 2 \text{ and } c_{t,0}[M, \Upsilon] = c_{t,2}[M, \Upsilon]\}$$

(If there is no such $t \in \mathbb{N}$ we let it be ∞ (i.e. ω for set theorists), we could also use “undefined”).

$$M \models_\iota \theta_{\Upsilon, \chi} \text{ iff } N_t[M, \Upsilon] \models \chi \text{ for } t = t_\iota[M, \Upsilon].$$

Case 2: For $\iota = 2$.

$$t_\iota[M, \Upsilon, \mathbf{t}] = \text{Min}\{t : \|N_{t+1}[M, \Upsilon]\| + (t+1) > \mathbf{t}(\|M\|) \text{ or } c_{t,0}[M, \Upsilon] = c_{t,1}[M, \Upsilon]\} \\ \text{and}$$

- (a) if $t = t_\iota[M, \Upsilon, \mathbf{t}] < \mathbf{t}(\|M\|)$, then the truth value of $\theta_{\Upsilon, \chi, \mathbf{t}}$ in M is true or false iff $N_t[M, \Upsilon] \models \chi$ or $N_t[M, \Upsilon] \models \neg\chi$ respectively and we write $M \models_\iota \theta_{\Upsilon, \chi, \mathbf{t}}$ or $M \models_\iota \neg\theta_{\Upsilon, \chi, \mathbf{t}}$ respectively.
- (b) if $t_\iota[M, \Upsilon, \mathbf{t}] = \mathbf{t}(\|M\|)$ we say “ $M \models_\iota \theta_{\Upsilon, \chi, \mathbf{t}}$ ” is undefined and we say “the truth value of $\theta_{\Upsilon, \chi, \mathbf{t}}$ in M ” is undefined.

Case 3: $\iota = 3$.

We restrict ourselves to standard Υ and let

$$t_\iota[M, \Upsilon, \mathbf{t}] = \text{Min}\{t : \|N_{t+1}[M, \Upsilon]\| > \mathbf{t}(\|M\|)\} \text{ and define } \models_\iota \text{ as in Case 2.}$$

Case 4: $\iota = 4$.

We restrict ourselves to standard Υ and

$$t_\iota[M, \Upsilon, \mathbf{t}] = \sup\{t : \text{for some } k < m_0 \text{ we have } \mathcal{P}_{t+1, k}[M, \Upsilon] \\ \text{has } > \mathbf{t}(\|M\|) \text{ members}\}$$

(so it can be ∞ ; i.e. ω for set theorists, but we can guarantee $\mathcal{P}_{t,0} = \{0, \dots, t-1\}$ so that this never happens) and define $\models_\iota$ as in Case 2.

We can replace in clause (b), $c_0 = c_1$ (i.e. $(\forall x)(P_0(x) \equiv P_1(x)) \ \& \ (\exists x)P_0(x)$) by a sentence χ .

1.3 Discussion: 1) The most smooth variant for our purpose is where $\mathcal{L}^* = \mathcal{L}_{\text{card}, \mathbf{T}}$ or $\mathcal{L}^* = L_{\text{card}, \mathbf{T}}$ and $\iota = 4$. From considering the motivation the most natural $\mathbf{T}$ is $\{n^m : m < \omega\}$, and $\iota = 3$.

2) For $\iota = 1, 2, 3$ some properties of M can be “incidentally” expressed by the logic, as the stopping time gives us some information on concerning cardinality can be expressed. This suggests preferring the option $\models_\iota$ undefined in case (b) rather than false.

3) If you like set theory, you can let t be any ordinal; but this is a side issue here; see §4.

Implicit in 1.2 (and alternative to 1.2) is

1.4 Definition. Let M, Υ be as in Definition 1.2 be given.

1) We say $(N, \bar{c})$ is an M -candidate or (M, Υ) -candidate

- (a) N is a transitive submodel of $\bigcup_t V_t[M]$
- (b) $\bar{c} = \langle c_\ell : \ell < m_1 \rangle, c_\ell \in N$ (or is undefined, so is really a unary relation on N of cardinality ≤ 1).

2) We say $(N', \bar{c}')$ is the $(\Upsilon, \mathfrak{t})$ -successor of $(N, \bar{c})$ if N', c'_ℓ are defined as in Definition 1.2, but

$$|N'| = |N| \cup \bigcup_{\ell < m_1} \mathcal{P}_\ell[N, \Upsilon]$$

$\mathcal{P}_\ell[N, \Upsilon] = A_\ell$ is $\{\{a : (N, \bar{c}) \models \psi_\ell(a, \bar{b})\} : \bar{b} \in {}^{\ell g(\bar{g}_\ell)} N\}$ if this family has $\leq \mathfrak{t}(M)$ members. A_ℓ is empty otherwise.

3) We define $N_t[M, \Upsilon, \mathfrak{t}]$ and $\bar{c}_t[M, \Upsilon, \mathfrak{t}]$ by induction on t as follows:

- for $t = 0$ it is M ($c_{t,\ell}$ undefined)
- for $t + 1, (N_{t+1}, \bar{c}_{t+1})$ is the $(\Upsilon, \mathfrak{t})$ - successor of $(N_t, \bar{c}_t)$, see below.

1.5 Claim. 1) if $(N, \bar{c})$ is an (M, Υ) -candidate, it has at most one $(\Upsilon, \mathfrak{t})$ -successor.
 2) The various definition fits, so we can use 1.4.

1.6 Discussion: How do we translate between the definitions above and [BGSh 533]?

- (i) an infinite structure I there corresponds to a τ -model here
- (ii) a state A there corresponds to a model of the form $N = N_t[M, \Upsilon]$ in 1.2 and $(N, \bar{c})$ in 1.4
- (iii) dynamic function there corresponds to $\mathcal{P}_\ell$ (that is $c_{t,\ell}$) here
- (iv) an object x is active at A in 5.1 there it corresponds to $x \in N$
- (v) a program 4.6 there corresponds to an Υ in 1.1(2) here (mainly the first order formulas used);
- (vi) the counting function in 4.7 there corresponds to the cardinality quantifier (1.1(6)) here
- (vii) the polynomial functions p, q in 5.1 there corresponds to $\mathfrak{t} \in T$ here.

Note that the $c_{t+1,\ell}$ can in the usual set theory manner be actually 7-place function from N_t to N_t or 7-place relation on N_t , or be the universe of N_t . Understanding this to interpret the successor step there to here we need that all parts of the program are expressible in $\mathcal{L}_{f.o.}$ (or $\mathcal{L}_{card}$). For the other direction we need to show f.o. operations can be expressed by the programs of ASM there (see 6.1 there), no problem (and not needed to show our results solved problems there).

§2 THE GENERAL ELIMINATION OF QUANTIFIERS
AND PROOFS IT'S NON-EXPRESSING

In Blass Gurevich Shelah [BGSh 533] we deal with the case of equality and permutations here we are using partial isomorphisms. It seems a reasonably precise way, so we shall later, hopefully, get a kind of inverse.

2.1 Discussion: Our aim is to have a family $\mathcal{F}$ of partial automorphisms as in Ehrenfeucht Fraisse games (or Karp), of the model M we analyze, not total automorphism (as in [BGSh 533]) - too restrictive. But it has to be lifted to the N_t 's. But their domains (and ranges) can contain an element of high rank. So we should not lose anything when we get up on t . The solution is $I \subseteq \{A : A \subseteq M\}$ closed downward and $\mathcal{F}$ (really $\langle \mathcal{F}_i : i < m_1 \rangle$), a family of partial automorphisms of M . So every $x \in N$ will have a support $A \in I$ and for $f \in \mathcal{F}$, its action on A determines its action on x , ($G(f)(x)$ in the section notation). It is not unreasonable to demand that there is a minimal one, still it is somewhat restrictive (or we have to add imaginary elements as in [Sh:a] or [Sh:c], not a very appetizing choice). But how come we in stage $t + 1$ succeed to add “all sets $X = X_{i,h}$ ” definable by $\psi_1(x, \bar{b}), \bar{b} \in {}^{\ell g(\bar{b})}N_t$?

The parameter $b_1, \dots, b_m$ each has a support say $A_1, \dots, A_m$ all in I , so we have enough mappings in the family, the new set has in a sense support $A = \bigcup_{\ell=1}^m A_\ell$, in the sense that suitable partial mappings do, if y has support B (BRy in this section notation) $A \cup B \subseteq \text{Dom}(f), f \upharpoonright A = \text{id}_A$ the mapping f induces in N , map y to a member of B .

But we are not allowed to increase the possible support and A though a kind of support is probably too large: I is not closed under union. But, if we add $X = X_{i,\bar{b}}$ we have to add all similar $X' = X_{i,\bar{b}'}$. So our strategy is to say no to looking for a support $A' \in I$. So fixing A' we like that if $f \in \mathcal{F}, f \upharpoonright A' = \text{id}_{A'}, A \subseteq \text{Dom}(f)$, then $f \upharpoonright A$ induces a mapping of $X_{i,\bar{b}}$ to some $X_{i,\bar{b}'}$, which we like to demand that will be equal thus justifying the statement “ A' supports X .”

How? We use our bound on the size of the computation. So we need a dichotomy: either there is such $A' \in I$ or the number $X_{i,b'}$ defined by $\psi_i(x, \bar{b}')$ is too large!! On this dichotomy hangs the proof.

However, we do not like to state this as a condition on N_t rather on M . We do not “know” how $\psi_1(x, \bar{b}')$ will act but for any possible A' this induces an equivalence relation on the images of A ($\mathcal{F}$ has to be large enough).

Actually, we can ignore f and develop set theory of elements demanding support in $\mathcal{F}$. So we break the proof to definition and claims.

We consider three variants of the logic: usual variant to make preservation clear, and the case with the cardinality quantifier.

We could use one $\mathcal{F}$ but we use $\langle \mathcal{F}_\ell : \ell \leq m_{\text{qd}}(*) \rangle$. Actually, for much of the treatment only $\mathcal{F}_0$ count.

Discussion: Each $a \in N$ will have support $A \in I$. Now should we in $(N, \bar{c})$ add the support of each $\bar{c}_k$ or this will be included? No! The $\bar{c}_t$ will have support $\emptyset$.

2.2 The Main Definition. 1) We say $\mathcal{Y} = (M, I, \mathcal{F})$ is a k -system if $\bar{m}^* = (m_{\text{qd}}(*), m_{\text{fv}}(*))$ and:

- (A) I is a family of subsets of $|M|$ (the universe of M) closed under subsets and each singleton belongs to it
[hint: intended as first approximation to the possible supports of the partial automorphism of M , the model of course; the intention is that M is finite]
Let $I[m] = \{\bigcup_{\ell=1}^m A_\ell : A_\ell \in I\}$
- (B) $\mathcal{F}$ is a family of partial automorphisms of M such that $A \subseteq \text{Dom}(f)$ & $A \in I \Rightarrow f''(A) \in I$; $\mathcal{F}$ closed under inverse (i.e. $f \in \mathcal{F}_\ell \Rightarrow f^{-1} \in \mathcal{F}_\ell$) and composition and restriction
- (C) if $f \in \mathcal{F}$ then $\text{Dom}(f)$ is the union of $\leq k$ members of I
- (D) if $f \in \mathcal{F}$ and $A_1, \dots, A_{k-1}, A_k \in I$ and $\ell \in \{1, \dots, k-1\} \Rightarrow A_\ell \subseteq \text{Dom}(f)$, then for some $g \in \mathcal{F}$ we have

$$f \upharpoonright \bigcup_{\ell=1}^{k-1} A_\ell \subseteq g$$

$$A_k \subseteq \text{Dom}(g)$$

- (E) if (α) then $(\beta)_1 \vee (\beta)_2$ where
 - (α) h is a function from some $[m] = \{1, \dots, m\}$, $\text{Rang}(h)$ belongs to $I[s]$, E is an equivalence relation on $H_h = \{h' : \text{for some } f \in \mathcal{F}_0, \text{Rang}(h) \subseteq \text{Dom}(f) \text{ and } h' = f \circ h\}$ satisfying
 - (*) if $h_1, h_2, h_3, h_4 \in H_h$ and there is $f \in \mathcal{F}$ such that $(\text{Rang } h_1) \cup (\text{Rang } h_3) \subseteq \text{Dom}(f)$ and $f \circ h_1 = h_3, f \circ h_2 = h_4$ then $(h_1 E h_2 \Leftrightarrow h_3 E h_4)$
 - (β)₁ there is $u \subseteq [m]$ such that $\text{Rang}(h \upharpoonright u) \in I$ and $(\forall h_1, h_2 \in H_h)(h_1 \upharpoonright u = h_2 \upharpoonright u \rightarrow h_1 E h_2)$
 - (β)₂ the number of E -equivalence classes is $> \mathfrak{t}(M)$
[hint: this is how from “there is some not too large number” we get “there is a support in I ”].

Note that without loss of generality h is one-to-one.

2) We say $(M, I, \mathcal{F})$ is a $(\mathfrak{t}, \bar{m})$ -system if $\bar{m} = (k, s)$ and it is a $(\mathfrak{t}, s)$ -dichotomical k -system; we may also say $(\mathfrak{t}, k, s)$ -system.

2.3 Definition. 1) We say $\mathcal{Y} = (M, I, \mathcal{F})$ is a super s -system if it is an $\bar{m}^*$ -system and in addition

(E)⁺ In clause (E), fixing some $A^* \in I[s]$ as a set of parameters, the number of equivalence classes is preserved; that is, if for $\ell = 1, 2$ we have $A_\ell \in I[s]$ and for some m and $h_\ell : [m] \rightarrow A$ with $\text{range} \in I$ and $\mathcal{E}_{\mathcal{Y}}(A_\ell, h_\ell)$ (see below) and $f \in \mathcal{F}$ such that $A_1 \cup \text{Rang}(h_1) \subseteq \text{Dom}(f)$ we have: f maps A_1 into A_2 , $f \circ h_1 = h_2$ and f maps E_1 to E_2 in the natural way, then the number of E_1 -equivalence classes is equal to the number of E_2 -equivalence classes.

2) Let $\mathcal{E}_{\mathcal{Y}}(A, h)$ be defined for $A \in I[s]$ and h a function from some $[m]$ into M with $\text{Rang}(h)$ belonging to I , $A \subseteq \text{Rang}(h)$ as follows:
 $\mathcal{E}_{\mathcal{Y}}(A, h)$ is the set of equivalence relations E on

$$H_{A,h} = \{h' : \text{for some } f \in \mathcal{F}, f \upharpoonright A = \text{id}, \text{Rang}(h) \subseteq \text{Dom}(f) \text{ and } h' = f \circ h\}.$$

such that the parallel (*) of clause (E) of Definition 2.2(2) holds. (So H_h is replaced by $H_{h,A}$, $f \upharpoonright A$ is the identity).

Let $\mathcal{E}_{\mathcal{Y}}(A) = \bigcup_h \mathcal{E}_{\mathcal{Y}}(A, h)$.

3) $\mathcal{Y} = (M, I, \mathcal{F})$ is a super $(\mathbf{t}, (k, s))$ -system if it is a super k -system and is $(\mathbf{t}, s)$ -dichotomical.

2.4 Observation: If $f \in \mathcal{F}$ maps A_1 to A_2 , then it induces a natural extension $\hat{f}$ of f mapping also $\mathcal{E}_{\mathcal{Y}}(A_1)$ onto $\mathcal{E}_{\mathcal{Y}}(A_2)$.

2.5 Definition. 1) Let $\mathcal{Y} = (M, I, \mathcal{F})$ be an k -system, M a τ -model, Υ is an inductive scheme for $\mathcal{L}_{f_0}(\tau^+)$ and $\bar{m}^* = \bar{m}(\Upsilon)$.

We say that $\mathcal{Z} = (N, \bar{c}, G, R) = (N^{\mathcal{Z}}, \bar{c}^{\mathcal{Z}}, G^{\mathcal{Z}}, R^{\mathcal{Z}})$ is a Υ -lifting of $\mathcal{Y}$ if

- (a) $(N, \bar{c})$ is an (M, Υ) -candidate so N is a transitive submodel of set theory with M as its set of urelements (see 1.4(1))
- (b) G is a function with domain $\mathcal{F}$
- (c) for $f \in \mathcal{F}$, $f \subseteq G(f)$ and $G(f)$ is a partial automorphism of N
- (d) if $f \in \mathcal{F}$, $g \in \mathcal{F}$, $f \subseteq g$ then $G(f) \subseteq G(g)$
- (e) R is a two-place relation written xRy such that
 $xRy \Rightarrow x \in I \ \& \ y \in N$
 [intention: x is a support of y]
- (f) if ARy and $f \in \mathcal{F}$, $A \subseteq \text{Dom}(f)$, then
 $y \in \text{Dom}(G(f))$ and $f \upharpoonright A = \text{id} \Rightarrow G(f)(y) = y$
- (g) $(\forall y \in N)(\exists A \in I)ARy$
- (h) if $A \in I$ and $A \subseteq \text{Dom}(f)$, $y \in \text{Dom}(G(f))$ then
 $ARy \Leftrightarrow f''(A)R(G(f)(y))$
- (i) for $f \in \mathcal{F}$, $G(f^{-1}) = (G(f))^{-1}$
- (j) for $f_1, f_2 \in \mathcal{F}$, $f = f_2 \circ f_1$ we have $G(f) \subseteq G(f_2) \circ G(f_1)$
- (k) if $c_\ell \in \text{Dom}(f)$ and $f \in \mathcal{F}$ then $f(c_\ell) = c_\ell$.

2.6 Definition. For $\mathcal{Y} = (M, I, \mathcal{F})$, a k -system the $0 - \Upsilon$ -lifting is $(M, \bar{c}, G, R)$ where

- (a) G is the identity on $\mathcal{F}$
- (b) $ARy \Leftrightarrow A \in I \ \& \ y \in A$
- (c) each c_ℓ undefined.

2.7 Fact: The $0 - \Upsilon$ -lifting (in Definition 2.6) exists and is a lifting.

2.8 Definition. Let $\mathcal{Y} = (M, I, \mathcal{F})$ an k -system and $\mathcal{Z} = (N, \bar{c}, G, R)$ be a Υ -lifting.

1) We say X is good or $(\mathcal{Y}, \mathcal{Z})$ -good if

- (a) X a subset of N
- (b) for some $A \in I$ we have A supports X (for our $\mathcal{Y}$ and $\mathcal{Z}$) which means:
if $f \in \mathcal{F}, BRy$ (so $B \in I, y \in N$)
and $A \cup B \subseteq \text{Dom}(f)$, and $f \upharpoonright A = \text{id}_A$
then $y \in X \Leftrightarrow G(f)(y) \in X$
(note: $G(f)(y)$ is well defined by clause (g) of 2.5)
- (c) $X \not\subseteq N$.

2) Let $\mathcal{P} = \mathcal{P}_{\mathcal{Y}, \mathcal{Z}}$ be the family of good subsets of N , let $\mathcal{R}$ be the two-place relation defined by: $A\mathcal{R}X$ iff A supports X , i.e. (b) of part (1) holds.

3) For $f \in \mathcal{F}$ and for good X such that $A\mathcal{R}X, A \in I$ when $A \subseteq \text{Dom}(f)$ we let

- (α) $(G^+(f))(X) = \{y \in N : \text{for some } g \in \mathcal{F} \text{ and } y' \in X \text{ we have}$
 $f \upharpoonright A \subseteq g \text{ and } G(g)(y') = y\}$ as $k \geq 3$, if the result is good
- (β) $G^+(f) \upharpoonright N = f$.

We can prove this, see 2.9 below.

4) We define $E = E_{\mathcal{Y}, \mathcal{Z}}$ as the following relation: $X_1 E X_2$ iff X_1, X_2 are good subsets of $N^{\mathcal{Z}}$ and for some $f \in \mathcal{F}$ we have $(G^+(f))(X_1) = X_2$; this is an equivalence relation as $k \geq 2$.

5) $\mathcal{Z}' = (N', \bar{c}', G', R')$ is a successor of $\mathcal{Z}$ if both are $\bar{m}^*$ -systems and:

- (a) $N \subseteq N' \subseteq N \cup \mathcal{P}_{\mathcal{Y}, \mathcal{Z}}$
- (b) $X_1 E X_2, X_1 \in \mathcal{P}_{\mathcal{Y}, \mathcal{Z}}, X_2 \in \mathcal{P}_{\mathcal{Y}, \mathcal{Z}} \Rightarrow [X_1 \in N' \leftrightarrow X_2 \in N']$
- (c) $G'(f)$ is defined as $G^+(f)$ from part (3)
- (d) R' is $R \cup [\mathcal{R} \upharpoonright (I \times N')]$.

6) $\mathcal{Z}' = (N', \bar{c}', G', R')$ is a full $\mathbf{t}$ -successor of $\mathcal{Z}$ iff above

$N' = N \cup \{X \in \mathcal{P}_{\mathcal{Y}, \mathcal{Z}} : |X/E_{\mathcal{Y}, \mathcal{Z}}| \leq \mathbf{t}(M)\}$. If we omit $\mathbf{t}$ we mean $\mathbf{t}(N) = \infty$.

7) $\mathcal{Z}' = (N', \bar{c}', B', R')$ is the true $(\Upsilon, \mathbf{t})$ -successor of $\mathcal{Z}$ if above:

- (a) $(N, \bar{c})$ is the $(\Upsilon, \mathbf{t})$ -successor of $(N, \bar{c})$, (see Definition 1.4(2)).

We now prove that Definition 2.8 is O.K. The functions defined are functions with the right domain and rang. The E 's are equivalence relations. This is included in the proof of 2.9.

2.9 Claim. Assume $\mathcal{Y}$ is a $(\mathbf{t}, \bar{m}^*)$ -system (see Definition 2.2(2)) and $\mathcal{Y}, \mathcal{Z}, m^*$ are as in Definition 2.8.

- 1) The true successor $\mathcal{Z}'$ of $\mathcal{Z}$ if exists is a successor of $\mathcal{Z}$.
- 2) A successor $\mathcal{Z}'$ of $\mathcal{Z}$ is a Υ -lifting of $(\mathcal{Y}, \bar{m}^*)$.
- 3) A full successor of $\mathcal{Z}$ exists.

Proof. 1) Trivial.

2) We check the clauses in Definition 2.5.

Clause (a): as N is transitive with M its set of urelements, and $X \in N' \setminus N \Rightarrow X \in \mathcal{P}_N \Rightarrow X \subseteq N$ also N' is transitive with M its set of urelements.

Clause (b): So we have defined G' above.

Clause (c): Let $f \in \mathcal{F}$, $G'(f) = f'$ and let $x, y \in N'$ belongs to the domain of f' and we should prove $N' \models y \in x \Leftrightarrow N' \models f'(y) \in f'(x)$ (we shall do more toward having clause (g) later).

If $x \in N$, then $f'(x)$ is necessarily in N , hence $N' \models y \in x \Rightarrow y \in N$ and $N' \models z \in f'(x) \Rightarrow z \in N$, so as $f' \upharpoonright N = f$ we are done. So we can assume $x \in N' \setminus N$, so x is a good subset of N , so for some $A_0 \in I$, $A_0 R x$. We define

$$z =: \{b \in N : \text{for some } g \in \mathcal{F} \text{ and } b' \in x \text{ we have} \\ f \upharpoonright A \subseteq g \text{ and } G(g)(b') = t\}.$$

We need the following

- (*)₁ z is a good subset of N with $A_1 = f''(A_0)$ a support of z
- (*)₂ $x = \{b' \in N : \text{for some } g \in \mathcal{F} \text{ and } b \in z \text{ we have} \\ f^{-1} \upharpoonright (f''(A_0)) \subseteq g \text{ and } G(g)(b) = b'\}$
- (*)₃ z does not belong to N
- (*)₄ $z = f'(x)$
- (*)₅ if B is another support of x , then $z' = z$ when $z' = \{b \in N : \text{for some } g \in \mathcal{F} \text{ and } a \in x \text{ we have } f \upharpoonright A \subseteq g \text{ and } G(g)(a) = b\}$.

Proof of ()₁.* Suppose $a, b \in N$ and $g_1 \in \mathcal{F}$ and $A_1 \subseteq \text{Dom}(g_1)$ and $g_1 \upharpoonright A_1$ is the identity and $a \in \text{Dom}[G(g_1)]$ and $b = G(g_1)(a)$. Now we should prove that $a \in z \Leftrightarrow b \in z$. It is enough to prove $\Rightarrow$ as applying it to g_1^{-1} we get the other implication. As $t = G(g_1)(s)$ necessarily by (i) of 2.5 for some support B_1 of s , $B_1 \subseteq \text{Dom}(g_1)$.

If $a \in z$ then by the definition of z we can find $g \in \mathcal{F}$ and $a' \in X$ such that $\text{id}_A \subseteq g$ and $G(g)(a') = a$. There is $B_2 \in I$ such that $B_2 R a'$ and $B_2 \subseteq \text{Dom}(g)$. As $A, B_1, B_2 \in I$ by clause (D) of Definition ? without loss of generality $B_1 \subseteq \text{Dom}(g)$

→ scite{2.1} undefined

(as $3 \leq m_{\text{qd}}(*)$, see Definition 1.1).

Let $g' = g_1 \circ g$, so $A \cup B_1 \subseteq \text{Dom}(g')$, $g' \upharpoonright A = g_1 \upharpoonright A = \text{id}_A$ hence $a' \in \text{Dom}(G(g'))$ and so $G(g')(a') = G(g_1)(G(g)(a')) = G(g_1)(a) = b$. (See Definition 2.5, clause (j).)

So g', a' witness $b \in z$ as required.

Proof of $()_2$.*

Similar.

Proof of $()_3$.* If $z \in N$ there is $A^* \in I$ such that $A^* R z$.

Now there is $f_1 \in \mathcal{F}$, $f \upharpoonright A \subseteq f_1$ such that $A^* \subseteq \text{Rang}(f_1)$, $f_1 \in \mathcal{F}$. So $z_1 = G(f_1^{-1})(z)$ is well defined and as in $(*)_2$ the proof of $(*)_1$ we can check that $\{b \in N : b \in z_1\} = x$; contradiction to “ $x \notin N$ ”.

Proof of $()_4$.*

Should be clear.

Proof of $()_5$.*

Should be clear.

Clause (d): Check.

Clauses (i),(j),(k): Check.

Clause (e): See Definition of R' .

Clause (f): Included in the poof of clause (d).

Clause (g): Check.

Clause (h):

The new case if: $A \subseteq \text{Dom}(f)$, $A \in I$, $X \in \text{Dom}(G'(f))$, $X \in N' \setminus N$. Then $AR'X \Leftrightarrow f''(A)R'(G'(f))(X)$ by clause (j) it is enough to prove the implication $\Rightarrow$.

There is $A_1 \in I$ such that $A_1 R' X$. If $\neg AR'X$ we can find $g \in \mathcal{F}$, $g \upharpoonright A = \text{id}_A$, and $z_0 \in \text{Dom } G(f)$, $z_1 = G(f)(z_0)$, such that $z_0 \in X \equiv z_1 \notin X$. We can find $B_0 \in I$ such that $A_0 R z_0$ and $B_1 \subseteq \text{Dom } g$ and let $A_1 =: g''(A_0)$. We can find $f_1, f \upharpoonright A \subseteq f_1, B_0 \subseteq \text{Dom}(f_1), f_1 \in \mathcal{F}$. Let $B'_0 = f_1(B_0)$.

Now chase arrows.

3) Straight. □_{2.9}

2.10 Claim. *Assume*

- (a) Υ is an inductive scheme for $\mathcal{L}_{\text{f.o.}}(\tau^+)$, $\bar{m}^* = \bar{m}^*(\Upsilon)$
- (b) k, s satisfy?
- (c) M is a finite τ -model, $\mathbf{t} \in \mathbf{T}$
- (d) $\mathcal{Y} = (M, I, \mathcal{F})$ is a $(\mathbf{t}, k, s)$ -system.

Then

- (α) if $N_t[M, \Upsilon, \mathbf{t}]$, $\bar{c}_t[M, \Upsilon, \mathbf{t}]$ are well defined, then for some $\mathcal{L} = (N, \bar{c}, R, G)$ a Υ -lifting we have $(N, \bar{c}, \mathcal{F})(N_t, \bar{c}_t)$.

2.11 Definition. 1) We say that $\mathcal{H}$ is a k -witness to the equivalence of $\mathcal{Y}_1$ and $\mathcal{Y}_2$ if

- (a) for $\ell = 1, 2$ we have $\mathcal{Y}_\ell = (M_\ell, I_\ell, \mathcal{F}_\ell)$ is a k -system
- (b) $\mathcal{G}$ is a family of partial isomorphisms from M_1 into M_2
- (c) for every $g \in \mathcal{H}$, we have $\text{Dom}(g) \in I_1$, $\text{Rang}(g) \in I_2$
- (d) if $g \in \mathcal{H}$ and $f_1 \in \mathcal{F}_1$ then $g \circ f_1 \in \mathcal{H}$
- (e) if $g \in \mathcal{H}$ and $f_2 \in \mathcal{F}_2$ then $f_2 \circ g \in \mathcal{H}$
- (f) if $g \in \mathcal{H}$ and $A \in I_1[k-1]$ and $B \in I_1$, then for some $g_1 \in \mathcal{H}$ we have $g \upharpoonright A \subseteq g_1$ and $B \subseteq \text{Dom}(g_1)$
- (g) if $g \in \mathcal{H}$ and $A \in I_2[k-1]$ and $B \in I_2$, then for some $g_1 \in \mathcal{H}$ we have $g^{-1} \upharpoonright A \subseteq g_1$ and $B \subseteq \text{Rang}(g_1)$.

2) We say that $\mathcal{H}$ is a (k, s) -witness to the equivalence of $(\mathcal{Y}_1, \mathbf{t}_1)$ and $(\mathbf{Y}_2, \mathbf{t}_2)$ if

- (i) $\mathcal{Y}_\ell$ is $(\mathbf{t}, k, s)$ -system, i.e. $(\mathbf{t}_\ell, s)$ -dichotomical k -system for $\ell = 1, 2$
- (ii) $\mathcal{H}$ is a witness to the k -equivalence of $\mathcal{Y}_1$ and $\mathcal{Y}_2$
- (iii) each $g \in \mathcal{G}$ preserved the possibility chosen in the definition or $(\mathbf{t}, s)$ -dichotomical.

3) We say that $\mathcal{H}$ is a super (k, s) -witness to the equivalence of $(\mathcal{Y}_1, \mathbf{t}_1)$ and $(\mathbf{Y}_2, \mathbf{t}_2)$ if

- (i) $\mathcal{Y}_\ell$ is a super $(\mathbf{t}_\ell, k, s)$ -system
- (ii) $\mathcal{H}$ is a witness to the k -equivalence of $\mathcal{Y}_1$ and $\mathcal{Y}_2$
- (iii) each $g \in \mathcal{H}$ preserve the cardinalities involved in the definition of super.

2.12 Main Conclusion: Assume

- (a) $\mathcal{Y}_\ell = (M_\ell, I_\ell, \mathcal{F}_\ell)$ is a $(\mathbf{t}, k, s)$ -system for $\ell = 1, 2$ and $\tau(M_\ell) = \tau$
- (b) $\mathcal{H}$ is a (k, s) -witness to the equivalence of $(\mathcal{Y}_1, \mathbf{t}_1)$ and $(\mathcal{Y}_2, \mathbf{t}_2)$
- (c) $\chi \in \mathcal{L}_{f.o.}(\tau^+)$, i.e. a first order sentence in the vocabulary $\tau^+ = \tau \cup \{\in\}$, and every subformula of χ has at most k -free variables
- (d) Υ is an inductive scheme not too complicated relative to k, s .

Then

- (α) the truth value of $\theta_{\Upsilon, \chi, \mathbf{t}_1}$ in M_1 and $\theta_{\Upsilon, \chi, \mathbf{t}_2}$ in M_2 are equal except possibly when: for some $\ell \in \{1, 2\}$ we have the truth value of $\theta_{\Upsilon, \chi, \mathbf{t}_\ell}$ in M_ℓ is undefined whereas that of $\theta_{\Upsilon, \chi, \mathbf{t}_{3-\ell}}$ in $M_{3-\ell}$ is well defined and $\mathbf{t}[M_\ell, \Upsilon, \mathbf{t}_\ell] < \mathbf{t}[M_{3-\ell}, \Upsilon, \mathbf{t}_{3-\ell}]$
- (β) for any t if $N^\ell = N_t[M_\ell, \Upsilon, \mathbf{t}_\ell]$ is well defined for $\ell = 1, 2$, then for every sentence χ such that every subformula has at most k -free variables, we have $N^1 \models \psi \Leftrightarrow N^2 \models \psi$.

Proof. Straight.

2.13 Claim. *In 2.10, 2.11 we can allow in the ψ_i and in the φ the cardinality quantifier provided that $\mathcal{V}$'s are super.*

Proof. Clearer than 2.10, 2.11.

2.14 Discussion We consider now some variants.

1) We can define a natural equivalence of two $\bar{m}^*$ -systems. Again the case with cardinality quantifiers is clearer.

This makes the proof of applications slightly different.

2) We have to consider the stopping times. If $\mathcal{L}^* = \mathcal{L}_{\text{car}, \mathbf{T}}$ or $\mathcal{L}_{\text{card}, \mathbf{T}}$ this is natural, (and they are stronger logics than the earlier variants). If we still would like to analyze in particular for the others, we should be careful how much information can be gotten by the time.

3) We can omit the c_ℓ 's if the models are rich enough by a first order formula. In N_{t+1} reconstruct the sequence $\langle (N_\ell, \bar{c}_\ell) : \ell \leq s \rangle$ (see §4).

§3 THE CANONICAL EXAMPLE

We apply §2 to the canonical example: random enough graph.

3.1 Definition. Let τ be a fixed vocabulary consisting of predicates only. We say M is a $(\mathbf{t}, k)$ -random τ -model if every quantifier free 1-type over $A \subseteq M, |A| < k$ (not explicitly inconsistent) is realized in M by at least $\mathbf{s}(\|M\|)$ elements. If $\mathbf{t} = \infty$ we may write k -random.

3.2 Definition. $\mathbf{T}_{\text{pol}}$ is $\{f_q : q \in \mathbb{Q}, q > 0\}$ where $f_q : \omega \rightarrow \omega$ is $f_q(n) = n^q$ or the least integer $\geq n^q$, more exactly.

3.3 Claim. *Assume*

- (a) M_ℓ is $(\mathbf{s}_\ell, k)$ -random τ -model for $\ell = 1, 2$
- (b) $3s \leq k$
- (c) $2^{(2^s) \times (\tau)} < \mathbf{t}_\ell(M_\ell) < \mathbf{s}_\ell(M_\ell) - s$
- (d) Υ is an inductive scheme for $\mathcal{L}_{\text{f.o.}}(\tau^+)$, χ a sentence in $\mathcal{L}_{\text{f.o.}}(\tau^+)$ and $m_{fv}(\Upsilon) \leq s$ and each subformula of any formula in Υ or χ has at most s free variables.

Then the truth values of $\theta_{\Upsilon, \chi, \mathbf{t}_1}$ in M_1 and $\theta_{\Upsilon, \chi, \mathbf{t}_2}$ in M_2 are equal except the case in 2.11.

Proof. Let $\ell = 1, 2$. We let $I_\ell = \{A \subseteq M_\ell : |A| \leq q\}$ and

$$\mathcal{F}_\ell = \{f : f \text{ is a partial automorphism of } M_\ell \\ \text{and } \text{Dom}(f) \text{ has at } \leq qk \text{ elements}\}$$

- (*)₁ $\mathcal{Y}_\ell = (M_\ell, I_\ell, \mathcal{F}_\ell)$ is a k -system
[why? the least obvious clause in Definition 2.2(1) is clause (D) which holds by Definition 3.1 above.]
- (*)₂ $\mathcal{Y}_\ell = (M_\ell, I_\ell, \mathcal{F}_\ell)$ is $(\mathbf{t}_\ell, s)$ -dichotomical
[why? let $m \in \mathbb{N}$ and let E be an equivalence relation on the set of $h : [m] \rightarrow M_\ell$ satisfying (*) of Definition 2.2(2). Without loss of generality h is one-to-one, so necessarily $m \leq s$. As $s \leq (qk)/2$ there is a quantifier free formula $\varphi(\bar{x}, \bar{y}) \in \mathcal{L}(\tau)$, $\ell g(\bar{x}) = \ell g(\bar{y}) = m$ such that $h_1 E h_2$ iff $M_1 \models \varphi[\langle h_1(i) : i \in [m] \rangle, \langle h_2(i) : i \in [m] \rangle]$.]

Can there be $\bar{a}_0, \bar{a}_1 \in {}^m(M_\ell)$ realizing the same quantifier free type (say $p(\bar{x})$) (over the empty set) which are not E -equivalent? If so we can find $\bar{a}_2 \in {}^m(M_\ell)$ realizing the same quantifier free type $p(\bar{x})$ and disjoint to $\bar{a}_0 \hat{\ } \bar{a}_1$ (use “ M_ℓ is $(3s)$ -random”), so without loss of generality $\bar{a}_0, \bar{a}_1$ are disjoint. Now we ask “are there disjoint $\bar{b}_0, \bar{b}_1 \in {}^m(M_1)$ realizing $p(\bar{x})$ which are E -equivalent? If yes, we easily get a contradiction to “ E an equivalence relation”. So easily there are at least $s(M_\ell) - m$ pairwise disjoint sequences realizing $p(\bar{x})$. Moreover, easily by transitivity for some

$u \subseteq [m]$ we have for $\bar{a}, \bar{b} \in {}^m(M_\ell)$ realizing $p(\bar{x})$, $M_\ell \models \varphi(\bar{a}, \bar{b})$ iff $\bar{a} \upharpoonright u = \bar{b} \upharpoonright u$ hence the number of equivalence classes is $\geq \mathbf{s}_\ell(\mathbf{M}_\ell) - m$ so we get one of the allowable answers in Definition 2.2(2).

So assume that there are no $\bar{a}_0, \bar{a}_1 \in {}^m(M_\ell)$ realizing the same quantifier free types over $\emptyset$, hence the number of E -equivalence classes is at most $|S_{qf}^m(\emptyset, M)| \leq 2^{(2^m) \times (\tau)}$, which is immaterial for us. but also each equivalence class is preserved by any $f \in \mathcal{F}$ which gives the other allowable answer in Definition 2.2(2).

Let

$$\mathcal{H} = \{f : f \text{ is a partial embedding of } M_1 \text{ into } M_2 \\ \text{with } \text{Dom}(f) \text{ with } \leq qk \text{ members}\}.$$

(*)₃ $\mathcal{H}$ is a (k, s) -witness to the equivalence of $(\mathcal{Y}_1, \mathbf{t}_1)$ and $(\mathbf{Y}_2, \mathbf{t}_2)$
[why? straight.]

So we can apply 2.11 and get the desired result. □_{3.3}

3.4 Conclusion: The logic $\mathcal{L}_4^t(\mathcal{L}_{f.o., \mathbf{T}})(\tau)$ satisfies the 0-1 law for finite random model with a fix probability for each predicate.

3.5 Comment: We can use time e.g. $\|M\|^{\log(\log \|M\|)}$, then in (*), if M has μ -membesr, then M has to be k_μ -random for appropriate k_μ .

3.6 Claim. Consider the vocabulary $\{P\}$, P unary. For every sentence $\psi \in \mathcal{L}_4^t(\mathcal{L}_{\text{car}, \mathbf{T}})(t)$ and say time $\mathbf{t}$, for any n large enough, if, $P_1, P_2 \subseteq [n]$, $|P_\ell|, |[n] \setminus P_\ell| \geq 2 \log_2(\mathbf{t}(M))$ then $([n], P_1) \models^{\mathbf{t}} \psi \Leftrightarrow ([n], P_2) \models^{\mathbf{t}} \psi$.

Proof. Here we can use automorphisms of $N_t[M, \mathbf{t}]$ as in [BGSh 533] or just use 3.3. □_{3.6}

3.7 Definition. 1) We say M is a τ -model with k -elimination of quantifiers if for every subset A_0, A_1 of M , $|A_0| = |A_1| < k$ and an isomorphism f from $M \upharpoonright A_0$ onto $M \upharpoonright A_1$ and $a_0 \in M$ there is $a_1 \in M$ such that $f = f \cup \{\langle a_0, a_1 \rangle\}$ is an isomorphism from $M \upharpoonright (A_0 \cup \{a_0\})$ onto $M \upharpoonright (A_1 \cup \{a_1\})$.

2) We replace “quantifiers” by “quantifier and counting” if we add: and the two sets $\{a'_0 \in M : a'_0, a_0 \text{ realize the same quantifier free type over } A_0\}$ and $\{a'_1 \in M \setminus a'_1, a_1 \text{ realize the same quantifier free type over } A_1\}$ has the same number of elements.

3.8 Claim. 1) We can in 3.3 weaken the demand “ M_ℓ is $(\mathbf{s}_\ell, k)$ -random τ -model” to

- (a) _{ℓ} ⁻ (α) M_ℓ has k -elimination of quantifier
- (β) if $\varphi(\bar{x}, \bar{y})$ is a quantifier free formula defining an equivalence relation and $\lg(\bar{x}) = \lg(\bar{y}) \leq s$ then the number of classes is $\geq \mathbf{t}_\ell(M_\ell)$ or each equivalence class is definable by a quantifier free type.

3.9 Remark. Parallel claims hold for the logic with the cardinality quantifier.

3.10 Claim. *Choiceless polynomial time + counting logic does not capture polynomial time.*

Proof. Use 2.12 on the question: $|P^M| \geq \|M\|/2$, similarly to 3.3 with $\tau = \{P\}$.

§4 CLOSING COMMENTS

We may consider

4.1 Definition. 1) A context is $(K, \mathcal{J})$ such that

- (a) K be a class of models with vocabulary (= the set of predicates) τ
- (b) $\mathcal{J}$ is a function such that
- (c) $\text{Dom}(\mathcal{J}) = K$
- (d) $\mathcal{J}(M)$ is a family of subsets of K , whose union is $|M|$, and closed under subsets
- (e) $\mathcal{J}$ is preserved by isomorphisms.

2) In 1) let

$\text{Seq}_{\mathcal{J}}^*(M) = \{\bar{a} : \bar{a} \text{ a sequence of members of } M \text{ of length } \alpha, \text{ Rang}(\bar{a}) \in \mathcal{J}(M)\}$.

3) We define a logic $\mathcal{L}$. For $k < \omega$ and $\alpha < \omega$ or just α an ordinal let us define the formulas in $\mathcal{L}_{k,\alpha}$ by induction on α , each formula φ has the form $\varphi(\bar{x}_0, \bar{x}_1, \dots, \bar{x}_{k-1})$, $k_1 \leq k$, where the $\bar{x}_\ell$'s are pairwise disjoint (finite) sequences of variables and every variable appearing freely in φ appear in one of those sequences (so any formula is coupled with such $\langle \bar{x}_0, \dots, \bar{x}_{k-1} \rangle$, probably some not actually appearing) (we may restrict to $\bar{x}_\ell$ finite).

$\alpha = 0$: quantifier free formula; i.e. any Boolean combination of atomic ones (with the right variables, of course).

$\alpha + 1$: α non-limit $\varphi(\bar{x}_0, \dots, \bar{x}_{k-1})$ is a Boolean combination of formulas of the form $(\exists \bar{y})\psi(\bar{x}_{i_0}, \dots, \bar{x}_{i_{k_2-2}}, \bar{y})$ where $k_2 \leq k$, $\psi(\bar{x}_{i_0}, \dots, \bar{x}_{i_{k_2-2}}, \bar{y}) \in \mathcal{L}_{k,\alpha}$.

α limit: $\mathcal{L}_{k,\alpha} = \bigcup_{\beta < \alpha} \mathcal{L}_{k,\beta}$.

$\alpha + 1, \alpha$ limit: $\mathcal{L}_{k,\alpha+1}$ is the set of Boolean combinations of members of $\mathcal{L}_{k,\alpha}$ of the right variables.

Let $\mathcal{L}_k = \bigcup_{\alpha} \mathcal{L}_{k,\alpha}$, $\mathcal{L}_* = \bigcup_{k < \omega} \mathcal{L}_k$, $\mathcal{L}_{k,<\alpha} = \bigcup_{\beta < \alpha} \mathcal{L}_{k,\beta}$.

4) We now define a satisfaction relation $M \models \varphi(\bar{a}_0, \dots, \bar{a}_{k-1})$ where $k_1 \leq k$ (depending on $\mathcal{J}$).

I.e. we define by induction on α , for $\varphi(\bar{x}_0, \dots, \bar{x}_{k-1}) \in \mathcal{L}_{k,\alpha}$, $\bar{a}_\ell \in \text{Seq}_{\mathcal{J}}^{\ell g(\bar{x}_\ell)}(M)$, when does $M \models \varphi[\bar{a}_0, \dots, \bar{a}_{k-1}]$ and when $M \models \neg \varphi[\bar{a}_0, \dots, \bar{a}_{k-1}]$. This is done naturally, in particular $M \models (\exists \bar{y})\varphi(\bar{a}_0, \dots, \bar{a}_{k_2-1}, \bar{y})$ iff for some $\bar{b} \in \text{Seq}_{\mathcal{J}}^{\ell g(\bar{y})}(M)$, (so $\text{Rang } \bar{b} \in \mathcal{J}(M)$) we have $M \models \varphi[\bar{a}_0, \dots, \bar{a}_{k_2-1}, \bar{b}]$.

4.2 Discussion: We may replace M by M^+ , adding elements coding each $A \in \mathcal{J}(M)$, with decoding by functions, still this requires infinitely many functions, we need to actually code any sequence listing each $A \in \mathcal{J}(M)$.

Still this framework seems to work more smoothly for its purposes.

4.3 Observation: In the framework of Definition 4.1, $M_1 \equiv_{\mathcal{L}_k} M_2$ iff there is a family $\mathcal{F}$ witnessing it which means

- (a) $f \in \mathcal{F} \Rightarrow f$ is a partial isomorphism from M_1 to M_2
- (b) $f \in \mathcal{F} \Rightarrow \text{Dom}(f)$ is the union of $\leq k$ members of $\mathcal{J}(M_1)$
- (c) $f \in \mathcal{F} \Rightarrow \text{Rang}(f)$ is the union of $\leq k$ members of $\mathcal{J}(M_2)$
- (d) if $f \in \mathcal{F}$, $A_1, \dots, A_{k-1}, A_k \in \mathcal{J}(M_1)$, $\ell < k \Rightarrow A_\ell \subseteq \text{Dom } f$, then for some

$$g \in \mathcal{F}, \bigcup_{\ell=1}^k A_\ell \subseteq \text{Dom}(g), g \upharpoonright \bigcup_{\ell=1}^{k-1} A_\ell \subseteq f$$
- (e) like (d) for f^{-1} , M_2, M_1 .

4.4 Discussion: 1) In §2 we can define $N_t[M] \equiv N_{\Upsilon, t}[M, \mathbf{t}]$ for every ordinal t , and so $V_\Upsilon[M, \mathbf{t}] = \cup \{N_{\Upsilon, \alpha}[M, \mathbf{t}] : \alpha \text{ an ordinal}\}$, see below. Now as in the case $i = 4$, the analysis in §2 works for this but it is not clear if we can get any interesting things.

Can this give interesting proofs of consistency for set theory with no choice but with urelement?

4.5 Definition. 1) We say Υ (from Definition 1.1) is pure if $m_1[\Upsilon] = 0$ so no c_ℓ .
 2) For pure Υ , let “ $\mathcal{Z}$ is the full $\mathbf{t}$ -successor of order t of $\mathcal{Z}$ ” as in 2.8 iterating t times, noting now the full $\mathbf{t}$ -successor is unique allowing t to be an ordinal and for limit ordinal t take just the union.

4.6 Fact: For any Υ we can find Υ' which is equivalent if we use in Definition 1.2 the case $i = 4$ (well when $\mathbf{t}(M_\ell)$ always is $\geq \beth$). In fact, we can reconstruct the sequence of $\langle c_{t, \ell} : t' < t \rangle$ in N_t .

4.7 The Main Definition. 1) We say $\mathcal{Y} = (M, I, \mathcal{F})$ is a k -system if $\bar{m}^* = (m_{\text{qd}}(*), m_{\text{fv}}(*))$ and:

- (A) I is a family of subsets of $|M|$ (the universe of M) closed under subsets and each singleton belongs to it
 [hint: intended as first approximation to the possible supports of the partial automorphism of M , the model of course; the intention is that M is finite]

$$\text{Let } I[m] = \left\{ \bigcup_{\ell=1}^m A_\ell : A_\ell \in I \right\}$$

- (B) $\mathcal{F}$ is a family of partial automorphisms of M such that $A \subseteq \text{Dom}(f)$ & $A \in I \Rightarrow f''(A) \in I$; $\mathcal{F}$ closed under inverse (i.e. $f \in \mathcal{F}_\ell \Rightarrow f^{-1} \in \mathcal{F}_\ell$) and composition and restriction

- (C) if $f \in \mathcal{F}$ then $\text{Dom}(f)$ is the union of $\leq k$ members of I
- (D) if $f \in \mathcal{F}$ and $A_1, \dots, A_{k-1}, A_k \in I$ and $\ell \in \{1, \dots, k-1\} \Rightarrow A_\ell \subseteq \text{Dom}(f)$, then for some $g \in \mathcal{F}$ we have

$$f \upharpoonright \bigcup_{\ell=1}^{k-1} A_\ell \subseteq g$$

$$A_k \subseteq \text{Dom}(g)$$

(E) if (α) then $(\beta)_1 \vee (\beta)_2$ where

(α) h is a function from some $[m] = \{1, \dots, m\}$, $\text{Rang}(h)$ belongs to $I[s]$, E is an equivalence relation on $H_h = \{h' : \text{for some } f \in \mathcal{F}_0, \text{Rang}(h) \subseteq \text{Dom}(f) \text{ and } h' = f \circ h\}$ satisfying

(*) if $h_1, h_2, h_3, h_4 \in H_h$ and there is $f \in \mathcal{F}$ such that $(\text{Rang } h_1) \cup (\text{Rang } h_3) \subseteq \text{Dom}(f)$ and $f \circ h_1 = h_3, f \circ h_2 = h_4$ then $(h_1 E h_2 \Leftrightarrow h_3 E h_4)$

(β)₁ there is $u \subseteq [m]$ such that

$\text{Rang}(h \upharpoonright u) \in I$ and

$(\forall h_1, h_2 \in H_h)(h_1 \upharpoonright u = h_2 \upharpoonright u \rightarrow h_1 E h_2)$

(β)₂ the number of E -equivalence classes is $> \mathbf{t}(M)$

[hint: this is how from “there is some not too large number” we get “there is a support in I ”].

We can connect this to §2 as follows.

4.8 Claim. Assume $\mathcal{Y}_0 = (M, I, \bar{\mathcal{F}})$, $\mathcal{L} = (N, \bar{c}, G)$, $\bar{m}^*$ are as in 2.2, 2.5. Then

(a) if $\varphi(\bar{x}) \in \mathcal{L}_{m_{\text{qf}}(*)}^\infty$ so $\ell g(\bar{x}) \leq m_{\text{qf}}(\bar{x})$, and every subformula of $\varphi(\bar{x})$ has $\leq m_{\text{qf}}(*)$ free variable, $\bar{a} \in {}^{\ell g(\bar{x})}N$, $\bigwedge_{\ell < \ell g(\bar{a})} A_\ell R a_\ell$ and $\bigcup_{\ell < \ell g(\bar{a})} A_\ell \subseteq \text{Dom}(f)$ and $f \in \mathcal{F}_0$ then

$$N \models \varphi[\dots, a_\ell, \dots] \equiv \varphi[\dots, G(f)(a_\ell), \dots]$$

(b) if $\varphi(\bar{x})$ has quantifier depth k and $\leq m_{\text{qf}}(*) + m_{\text{qd}}(*) - k$ free variables, $f \in \mathcal{F}_{m_{\text{qf}}(*)}$ and $\dots, a_\ell, \dots \in \text{Dom}(G(f))$ then

$$N \models \varphi[\dots, a_\ell, \dots] \equiv \varphi[\dots, G(f)(a_\ell), \dots].$$

Proof. Straight.

4.9 Conclusion 1) Assume $\mathcal{Y}$ is super (see 2.3). Then we can define R_t, G_t for every t (N_t the “computation” in time t) such that

$$(M, \bar{c}_0, G_0, R_0) \text{ is 0-lifting}$$

$$(N_{t+1}, \bar{c}_{t+1}, R_{t+1}, \langle G_{t+1} \rangle) \text{ is a lifting, successor of } (N_t, \bar{c}_t, G_t, R_t).$$

2) So the formula the $\bar{\varphi}$ defines is preserved by $f \in \mathcal{F}_0$.

Proof. Straight.

REFERENCES.

- [BGSh 533] Andreas Blass, Yuri Gurevich, and Saharon Shelah. Choiceless Polynomial Time. *Annals of Pure and Applied Logic*, **accepted**.
- [Sh:a] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam-New York, xvi+544 pp, \$62.25, 1978.
- [Sh:c] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, xxxiv+705 pp, 1990.